

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

| Basic Information | |
|--------------------|---|
| Product Name | Anti-STAT1 Antibody |
| Gene Name | STAT1 |
| Source | Rabbit |
| lsotype | IgG |
| Species Reactivity | human |
| Tested Application | WB, FCM, ELISA |
| Contents | 500 ug/ml antibody with PBS, 0.02% NaN3 , 1 mg BSA and 50% glycerol. |
| Immunogen | E.coli-derived human STAT1 recombinant protein (Position: K145-D744). |
| concentration | 500 ug/ml |
| Purification | Immunogen affinity purified. |
| Dilution Ratios | Western blot(WB): 1:500-2000 Flow cytometry (FCM):1-3 μ g/1x10 ⁶ cells ELISA: 1:100-1000 |

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Signal transducer and activator of transcription 1?(STAT1) is a?transcription factor?which in humans is encoded by the?STAT1?gene. The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described.

Selected Validation Data

Product datasheet Anti-STAT1 Antibody Catalog Number: A00036-3





Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Figure 1. Western blot analysis of anti- STAT1 antibody (A00036-3).The sample well of each lane was loaded with 30ug of sample under reducing conditions.Lane 1: human 293T whole cell lysates,Lane 2: human Hela whole cell lysates,Lane 3: human PC-3 whole cell lysates.

Use rabbit anti- STAT1 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog#EK1002). A specific band was detected for STAT1 at approximately 91KD. The expected band size for STAT1 is at 87KD.