

Basic Information

Product Name	Anti-HO-1/HMOX1 Antibody
Gene Name	HMOX1
Source	Rabbit
Isotype	IgG
Species Reactivity	human
Tested Application	WB, FCM, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived human Heme Oxygenase 1/HMOX1 recombinant protein (Position: E19-R269).
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	33KD
Dilution Ratios	Western blot (WB): 1:500-2000 Flow cytometry (FCM): 1-3 µg/1×10 ⁶ cells ELISA: 1:100-1000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

HMOX1 (heme oxygenase (decycling) 1), also known as HO-1, is a human gene that encodes for the enzyme heme oxygenase 1. It is an essential enzyme in heme catabolism, it cleaves heme to form biliverdin. HMOX1 belongs to the heme oxygenase family. The HMOX1 gene is located on the long (q) arm of chromosome 22 at position 12.3, from base pair 34,101,636 to base pair 34,114,748. HMOX1, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is subsequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitter. HMOX1 activity is induced by its substrate heme and by various nonheme substances.

Selected Validation Data




Figure 1. Western blot analysis of anti- HO-1/HMOX1 antibody (A00253-2). The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human A549 whole cell lysates,

Lane 3: human 293T whole cell lysates.

Use rabbit anti- HO-1/HMOX1 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002). A specific band was detected for HO-1/HMOX1 at approximately 33KD. The expected band size for HO-1/HMOX1 is at 33KD.