Special NO.1, International Enterprise Center,

2nd Guanshan Road, Wuhan, China **Web:** www.boster.com.cn **Phone:** +86 027-67845390 **Fax:** +86 027-67845390 **Email:** boster@boster.com.cn

Basic Information	
Product Name	Anti-MMP2 Antibody
Gene Name	MMP2
Source	Rabbit
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS $_{2}$ 0.02% NaN3 , 1 mg BSA and 50% glycerol.
Immunogen	E. coli-derived human MMP2 recombinant protein (Position: K429-C660). Human MMP2 shares 95.7% and 95.3% amino acid (aa) sequence identity with mouse and rat MMP2, respectively.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	72KD
Observed MW	72KD

Storage

Dilution Ratios

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

1:100-1000

Western blot(WB):1:500-2000

ELISA:

Background Information

Matrix metalloproteinase-2 (MMP2) is a Type IV collagenase, 72-kD, which is also known as gelatinase and is a member of a group of secreted zinc metalloproteases. The MMP2 gene is 17 kb long with 13 exons varying in size from 110 to 901 bp and 12 introns ranging from 175 to 4,350 bp, located within a region of human chromosome 16q13. In addition, the extra exons encode the amino acids of the fibronectin-like domain which has so far been found in only the 72- and 92-kDa type IV collagenase. MMP2, which has a critical role in the binding of progelatinase A and TIMP4 via the C-terminal hemopexin-like domain (C domain), is functionally associated on the surface of angiogenic blood vessels. Not only is a likely effector of endometrial menstrual breakdown, MMP2 is also effector and regulator of the inflammatory response. Moreover, MMP2 could be helpful in diagnosing Takayasu arteritis.

Reference

Anti-MMP2 Antibody被引用在9文献中。



BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Selected Validation Data

100KD — 70KD — -55KD — 35KD — 25KD — Figure 1. Western blot analysis of MMP2 using anti-MMP2 antibody (A00286). lane 1: recombinant human MMP2 protein 1ng probed with a goat anti-rabbit IgG-HRP secondary antibody . The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) A specific band was detected for MMP2 at approximately 72KD. The expected band size for MMP2 is at 74KD.