

Basic Information

Product Name	Anti-iNOS/NOS2 Antibody
Gene Name	NOS2
Source	Rabbit
Isotype	IgG
Species Reactivity	mouse
Tested Application	WB, FCM, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived mouse iNOS/Nos2 recombinant protein (Position: I40-L1099).
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	130KD
Dilution Ratios	Western blot(WB): 1:500-2000 Flow cytometry (FCM): 1-3 μ g/1 \times 10 ⁶ cells ELISA: 1:100-1000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Nitric oxide synthase, inducible is an enzyme that in humans is encoded by the NOS2 gene. Nitric oxide (NO) is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter; it is implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. Three different NOS isoforms have been identified which fall into two distinct types, constitutive and inducible. The inducible NOS (iNOS) isoform is expressed in a variety of cell types and tissues in response to inflammatory agents and cytokines. The human iNOS (NOS2) gene is approximately 37 kb in length and consists of 26 exons and 25 introns. NOS2-derived NO is a prerequisite for cytokine signaling and function in innate immunity.

Selected Validation Data




Figure 1. Western blot analysis of anti- NOS2 antibody (A00368-4).

The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: mouse RAW264.7(-LPS) whole cell lysates,

Lane 2: mouse RAW264.7(+LPS) whole cell lysates.

Use rabbit anti- NOS2 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an

Enhanced Chemiluminescent detection (ECL) kit (Catalog#EK1002).

A specific band was detected for NOS2 at approximately 130KD.

The expected band size for NOS2 is at 130KD.