

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Basic Information	
Product Name	Anti-TIMP1 Antibody
Gene Name	TIMP1
Source	Rabbit
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS $_{\odot}$ 0.02% NaN3 , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived human TIMP1 recombinant protein (Position: H30-S164).
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	23-28KD
Dilution Ratios	Western blot(WB):1:500-2000 (ELISA): 1:100-1000
	(ELISA): 1:100-1000

Storage

12 months from date of receipt, -20° C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

TIMP metallopeptidase inhibitor 1, also known as TIMP1, a tissue inhibitor of metalloproteinases, is a glycoprotein that is expressed from the several tissues of organisms. This protein is a member of the TIMP family. The glycoprotein is a natural inhibitor of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. TIMP was found to be located about 22 cM proximal to OTC (300461).

Reference

Anti-TIMP1 Antibody被引用在1文献中。

Selected Validation Data

Product datasheet Anti-TIMP1 Antibody

Catalog Number: A00561-1

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn



Figure 1. Western blot analysis of Timp-1 using anti-Timp-1 antibody (A00561-1). The sample well of each lane was loaded with

Lane 1: human SK-OV-3 whole cell lysates,

Lane 2: human 22RV1 whole cell lysates,

Lane 3: rat ovarian tissue lysates,

Lane 4: mouse ovarian tissue lysates.

Use rabbit anti- Timp-1 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) . A specific band was detected for Timp-1 at approximately 23-28KD. The expected band size for Timp-1 is at 23KD.