

## Basic Information

<b>Product Name</b>	Human 14-3-3 Epsilon/YWHAE Recombinant Protein	
<b>Gene Name</b>	YWHAE	
<b>Source</b>	Rabbit	
<b>Isotype</b>	IgG	
<b>Species Reactivity</b>	human, mouse, rat	
<b>Tested Application</b>	WB, IHC, ICC/IF, FCM, ELISA	
<b>Contents</b>	500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol.	
<b>Immunogen</b>	E.coli-derived human YWHAE recombinant protein (Position: M1-Q255).	
<b>concentration</b>	500 ug/ml	
<b>Purification</b>	Immunogen affinity purified.	
<b>Observed MW</b>	29KD	
<b>Dilution Ratios</b>	Western blot(WB):	1:500-2000
	Immunohistochemistry in paraffin section IHC	1:50-400
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400
	Flow cytometry (FCM):	1-3 $\mu$ g/1x10 <sup>6</sup> cells
	ELISA:	1:100-1000

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

## Background Information

14-3-3 protein epsilon is a protein that in humans is encoded by the YWHAE gene. This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the mouse ortholog. It interacts with CDC25 phosphatases, RAF1 and IRS1 proteins, suggesting its role in diverse biochemical activities related to signal transduction, such as cell division and regulation of insulin sensitivity. It has also been implicated in the pathogenesis of small cell lung cancer. Two transcript variants, one protein-coding and the other non-protein-coding, have been found for this gene.

## Selected Validation Data

# Human 14-3-3 Epsilon/YWHAE Recombinant Protein

Catalog Number: **A01687-4**

Web: [www.boster.com.cn](http://www.boster.com.cn) Phone: +86 027-67845390 Fax: +86 027-67845390 Email: [boster@boster.com.cn](mailto:boster@boster.com.cn)




Figure 1. Western blot analysis of anti-YWHAE antibody (A01687-4). The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human hela whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: human hepg2 whole cell lysates, Lane 4: human SH-SY5Y whole cell lysates, Lane 5: human HEK293 whole cell lysates, Lane 6: human SW620 whole cell lysates, Lane 7: human A549 whole cell lysates, Lane 8: human Raji whole cell lysates, Lane 9: Rat brain tissue lysates, Lane 10: Mouse brain tissue lysates, Lane 11: Mouse NIH/3T3 whole cell lysates, Lane 12: Mouse RAW264.7 whole cell lysates, Use rabbit anti-YWHAE 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002). A specific band was detected for YWHAE at approximately 29KD. The expected band size for YWHAE is at 29KD.