

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

ation
Anti-Cyclin D1/CCND1 Antibody
CCND1
Rabbit
lgG
human, mouse, rat
WB
500 ug/ml antibody with PBS $ ightarrow$ 0.02% NaN3 , 1 mg BSA and 50% glycerol.
A synthetic peptide corresponding to a sequence at the N-terminal of human Cyclin D1, different from the related mouse sequence by two amino acids.
500 ug/ml
Immunogen affinity purified.
34KD
Western blot(WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Cyclin D1, also known as CCND1, is a human gene. The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclin D1 encodes the regulatory subunit of a holoenzyme that phosphorylates and inactivates the retinoblastoma protein and promotes progression through the G1-S phase of the cell cycle. Amplification or overexpression of cyclin D1 plays pivotal roles in the development of a subset of human cancers including parathyroid adenoma, breast cancer, colon cancer, lymphoma, melanoma, and prostate cancer. The cyclin D1 gene is overexpressed in human breast cancers and is required for oncogene-induced tumorigenesis. Brisken et al.(2003) found that prolactin(PRL; 176760) induced IGF2(147470) mRNA and IGF2 induced cyclin D1 protein expression in mouse mammary epithelial cultures. And they also concluded that IGF2 is a mediator of prolactin-induced alveologenesis and that prolactin, IGF2, and cyclin D1 are components of a developmental pathway in mammary gland.

Reference

Anti-Cyclin D1/CCND1 Antibody 被引用在29文献中。

Product datasheet Anti-Cyclin D1/CCND1 Antibody Catalog Number: BA0770



2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Selected Validation Data