

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Basic Information	
Product Name	Anti-MMP9 Antibody
Gene Name	MMP9
Source	Rabbit
lsotype	lgG
Species Reactivity	mouse, rat
Tested Application	WB, IHC
Contents	500 ug/ml antibody with PBS $ ightarrow$ 0.02% NaN3 , 1 mg BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human MMP-9(689-705aa NQVDQVGYVTYDILQCP), different from the related mouse sequence by two amino acids, and from the related rat sequence by three amino acids.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	78KD
Dilution Ratios	Western blot(WB):1:500-2000Immunohistochemistry in paraffin section (IHC):1:50-400(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20mins is required for the staining of formalin/paraffin sections.) Optimal working dilutionsmust be determined by end user.

Storage

12 months from date of receipt, -20° C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Matrix metallopeptidase 9(MMP-9), also known as 92 kDa type IV collagenase, 92 kDa gelatinase or gelatinase B(GELB), is an enzyme that in humans is encoded by the MMP9 gene. Proteins of the matrix metalloproteinase(MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling.

Reference

Anti-MMP9 Antibody被引用在14文献中。



Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Selected Validation Data

