# Product datasheet Anti-ATG5 Antibody Catalog Number: BA3525-2



**BOSTER BIOLOGICAL TECHNOLOGY** 

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Basic Information	
Product Name	Anti-ATG5 Antibody
Gene Name	ATG5
Source	Rabbit
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS ,0.02% NaN3 , 1 mg BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human APG5L(82-97aa DRFDQFWAINRKLMEY), identical to the related mouse sequence, and different from the related rat sequence by one amino acid.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	55KD
Dilution Ratios	Western blot(WB):1:500-2000

## **Storage**

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

### **Background Information**

Autophagy protein 5 is a protein that in humans is encoded by the ATG5 gene. It is also known as APG5 or ASP, and this gene is mapped to 6q21. It is found that knockdown of ATG5 in hepatocytes increased triglyceride levels with oleate or a second endogenous stimulus for triglyceride formation. These hepatocytes with ATG5 knockdown also had increased lipid droplet number and size. ATG5 is an E3 ubiquitin ligase which is necessary for autophagy due to its role in autophagosome elongation. It is activated by ATG7 and forms a complex with ATG12 and ATG16L1. This complex is necessary for LC3-1 conjugation to PE to form LC3-II.

#### Reference

Anti-ATG5 Antibody被引用在6文献中。

**BOSTER BIOLOGICAL TECHNOLOGY** 

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

#### **Selected Validation Data**




Figure 1. Western blot analysis of Anti-ATG5 antibody (BA3525-2). The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: Panc-1 whole cell lysates, Lane 2: HT1080 whole cell lysates,

Lane 3: U251 whole cell lysates, Lane 4: C6 whole cell lysates,

Lane 5: mouse testis tissue lysates.

Use rabbit Anti-ATG5 1:1000, probed with a goat Anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002). A specific band was detected for ATG5 at approximately 55KD. The expected band size for ATG5 is at 32KD.