

Basic Information

Product Name	Anti-Lamin A/C Antibody	
Gene Name	LMNA	
Source	Rabbit	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Immunogen	A synthesized peptide derived from human Lamin A/C	
Purification	Affinity-chromatography	
Observed MW	70-74KD	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry in paraffin section (IHC):	1:20-100
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:20-100
	Immunoprecipitation:	1:20
	Flow cytometry (FCM):	1:20

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Lamins are structural protein components of the nuclear lamina, a protein network underlying the inner nuclear membrane that determines nuclear shape and size. There are three types of lamins, A, B and C. The lamin A/C (LMNA) gene contains 12 exons. Alternative splicing within exon 10 gives rise to two different mRNAs that code for pre-lamin A and lamin C. Lamin A/C is mapped to 1q21.2-q21.3 and mutations in this gene cause a variety of human diseases including Emery-Dreifuss muscular dystrophy, dilated cardiomyopathy, and Hutchinson-Gilford progeria syndrome. Lamin A/C deficiency is thus associated with both defective nuclear mechanics and impaired mechanically activated gene transcription.

Selected Validation Data

Product datasheet

Anti-Lamin A/C Antibody

Catalog Number: **BM4105**


BOSTER

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center,
2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn **Phone:** +86 027-67845390 **Fax:** +86 027-67845390 **Email:** boster@boster.com.cn



Western blot analysis of Lamin A/C expression in HeLa whole cell lysates.