

Basic Information

Product Name	Anti-Histone H3 (mono+di+tri methyl K14) Antibody
Gene Name	H3C1/H3C2/H3C3/H3C4/H3C6/H3C7/H3C8/H3C10/H3C11/H3C12
Source	Rabbit
Isotype	IgG
Species Reactivity	human,mouse
Tested Application	WB
Contents	500 ug/ml;Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Immunogen	A synthesized peptide derived from human Histone H3 (mono+di+tri methyl K14)
Purification	Affinity-chromatography
Observed MW	15KD
Dilution Ratios	Western blot (WB): 1:400-800 Immunocytochemistry/Immunofluorescence(ICC/IF):1:20-100

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases.

Selected Validation Data

Product datasheet

Anti-Histone H3 (mono+di+tri methyl K14) Antibody

Catalog Number: **BM4337**


BOSTER

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center,
2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn



Western blot analysis of Histone H3 (mono+di+tri methyl K14) expression in HeLa cell lysate.