

Basic Information

Product Name	Anti-Cyclin B1/CCNB1 Antibody		
Gene Name	CCNB1		
Source	Rabbit		
Isotype	IgG		
Species Reactivity	human, mouse		
Tested Application	WB, IHC, ICC/IF, FCM		
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		
Immunogen	A synthesized peptide derived from human Cyclin B1		
concentration	500 ug/ml		
Purification	Affinity-chromatography		
Observed MW	55KD		
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry in paraffin section (IHC): 1:20-100		

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

CCNB also known as Cyclin B1, is a protein that in humans is encoded by the CCNB1 gene, and it is mapped to 5q13.2. The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. CCNB contributes to the switch-like all or none behavior of the cell in deciding to commit to mitosis. Its activation is well-regulated, and positive feedback loops ensure that once the cyclin B1-Cdk1 complex is activated, it is not deactivated.

Selected Validation Data

Western blot analysis of Cyclin B1 expression in HeLa cell lysate.

