

## Basic Information

<b>Product Name</b>	Anti-Caspase 9/CASP9 Antibody
<b>Gene Name</b>	CASP9
<b>Source</b>	Rabbit
<b>Isotype</b>	IgG
<b>Species Reactivity</b>	human, mouse
<b>Tested Application</b>	WB, IP
<b>Contents</b>	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Immunogen</b>	A synthesized peptide derived from human cleaved Caspase-9
<b>concentration</b>	500 ug/ml
<b>Purification</b>	Affinity-chromatography
<b>Observed MW</b>	46,39,37,35KD
<b>Dilution Ratios</b>	Western blot (WB): 1:500-2000 Immunoprecipitation:1:20

## Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


## Background Information

CASP9 is also known as MCH6 or APAF3. This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein can undergo autoproteolytic processing and activation by the apoptosome, a protein complex of cytochrome c and the apoptotic peptidase activating factor 1; this step is thought to be one of the earliest in the caspase activation cascade. This protein is thought to play a central role in apoptosis and to be a tumor suppressor. Alternative splicing results in multiple transcript variants.

## Reference

Anti-Caspase 9/CASP9 Antibody被引用在2文献中。

## Selected Validation Data



Western blot analysis of cleaved Caspase-9 Antibody expression in HeLa cell lysate treated with staurosporine.