

Basic Information

Product Name	Anti-SOD1 Antibody	
Gene Name	SOD1	
Source	Rabbit	
Isotype	IgG	
Species Reactivity	human, rat	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Immunogen	A synthesized peptide derived from human SOD1	
concentration	500 ug/ml	
Purification	Affinity-chromatography	
Observed MW	18KD	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry in paraffin section (IHC): 1:20-100	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Superoxide dismutases (SOD) are a class of enzymes that catalyze the dismutation of superoxide into oxygen and hydrogen peroxide. As such, they are an important antioxidant defense in nearly all cells exposed to oxygen. One of the exceedingly rare exceptions is *Lactobacillus plantarum* and related lactobacilli, which use a different mechanism. Cu,Zn-SOD was found widely distributed in the cell cytosol and in the cell nucleus, consistent with it being a soluble cytosolic protein. Mitochondria and secretory compartments did not label for this protein. In human cells, peroxisomes showed a labeling density slightly less than that of cytoplasm.

Reference

Anti-SOD1 Antibody被引用在1文献中。

Selected Validation Data

Western blot analysis of CD80 expression in Jurkat cell lysate.

