

Basic Information

Product Name	BrdU
Gene Name	
Source	
Species Reactivity	化学试剂
Tested Application	化学试剂,配合BrdU抗体使用
Contents	
Immunogen	
Purification	
Observed MW	
Dilution Ratios	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution.
Avoid repeated freezing and thawing

Background Information

Reference

BrdU被引用在4文献中。

Roxarsone induces angiogenesis via PI3K/Akt signaling Wang Y, Yin D, Xu C, Wang K, Zheng L, Zhang Y. Cell Biosci. 2016 Sep 28;6:54. eCollection 2016.	PubMed:27708768
Melatonin promotes diabetic wound healing in vitro by regulating keratinocyte activity Song R, Ren L, Ma H, Hu R, Gao H, Wang L, Chen X, Zhao Z, Liu J. Am J Transl Res. 2016 Nov 15;8(11):4682-4693. eCollection 2016.	PubMed:27904671
Exogenous Nkx2.5- or GATA-4-transfected rabbit bone marrow mesenchymal stem cells and myocardial cell co-culture on the treatment of myocardial infarction in rabbits Li P, Zhang L Mol Med Rep. 2015 Aug;12(2):2607-21. doi: 10.3892/mmr.2015.3775. Epub 2015 May 12.	PubMed:25975979
Optimal concentration and time window for proliferation and differentiation of neural stem cells from embryonic cerebral cortex: 5% oxygen preconditioning for 72 hours Yuan LL, Guan YJ, Ma DD, Du HM Neural Regen Res. 2015 Sep;10(9):1516-22. doi: 10.4103/1673-5374.165526	PubMed:26604915

Product datasheet

BrdU

Catalog Number: ED1100

BOSTER
antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY
Special NO.1, International Enterprise Center, 2nd
Guanshan Road, Wuhan, China

Web: www.boster.com.cn *Phone:* +86 27 67845390 *Fax:* +86 27 67845390 *Email:* boster@boster.com.cn

Selected Validation Data

