# Anti-FOXO3A/FOXO3 Antibody (Clone#OTI5F11)

Catalog Number: MA00252



**BOSTER BIOLOGICAL TECHNOLOGY** 

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Basic Information	
Product Name	Anti-FOXO3A/FOXO3 Antibody (Clone#OTI5F11)
Gene Name	FOXO3
Source	Mouse
Isotype	lgG1
Species Reactivity	human, mouse, rat
Tested Application	WB, IHC
Contents	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Immunogen	Human recombinant protein fragment corresponding to amino acids 374-673 of human FOXO3 (NP_001446) produced in E.coli.
concentration	500 ug/ml
Purification	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Dilution Ratios	Western blot (WB): 1:2000 Immunohistochemistry in paraffin section (IHC):1:150

#### **Storage**

Stable for 12 months from date of receipt. Store at -20°C as received.

### **Background Information**

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. This gene likely functions as a trigger for apoptosis through expression of genes necessary for cell death. Translocation of this gene with the MLL gene is associated with secondary acute leukemia. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq, Jul 2008]

### **Selected Validation Data**

#### **Product datasheet**

## Anti-FOXO3A/FOXO3 Antibody (Clone#OTI5F11)

Catalog Number: MA00252



**BOSTER BIOLOGICAL TECHNOLOGY** 

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn




Figure 1. Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-FOXO3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, MA00252) (1:150)