

Basic Information

Product Name	Anti-c-Met/MET Antibody
Gene Name	MET
Source	Rabbit
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS , 0.02% Na ₂ S ₂ O ₃ , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived human Met recombinant protein (Position: D208-S407). Human Met shares 90% and 91% amino acid (aa) sequences identity with mouse and rat Met, respectively.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	156KD
Dilution Ratios	Western blot(WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

c-Met, also called MET and hepatocyte growth factor receptor (HGFR), is a protein that in humans is encoded by the MET gene. It is mapped to 7q31.2. The protein possesses tyrosine kinase activity. MET is a membrane receptor that is essential for embryonic development and wound healing. It induces several biological responses that collectively give rise to a program known as invasive growth. MET is deregulated in many types of human malignancies, including cancers of kidney, liver, stomach, breast, and brain. Normally, only stem cells and progenitor cells express MET, which allows these cells to grow invasively in order to generate new tissues in an embryo or regenerate damaged tissues in an adult. However, cancer stem cells are thought to hijack the ability of normal stem cells to express MET, and thus become the cause of cancer persistence and spread to other sites in the body.

Selected Validation Data




Figure 1. Western blot analysis of Met (c-Met) using anti-Met (c-Met) antibody (PB0146). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. lane 1: rat liver tissue lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Met (c-Met) antigen affinity purified polyclonal antibody (Catalog # PB0146) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Met (c-Met) at approximately 154KD. The expected band size for Met (c-Met) is at 154KD.