

Basic Information

Product Name	Anti-PI3 Kinase p110 Alpha/PIK3CA Antibody
Gene Name	PIK3CA
Source	Rabbit
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS , 0.02% Na ₂ S ₂ O ₃ , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived human PIK3CA recombinant protein (Position: H936-N1068). Human PIK3CA shares 98% amino acid (aa) sequence identity with mouse PIK3CA.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	110KD
Dilution Ratios	Western blot(WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Phosphatidylinositol-4,5-bisphosphate 3-kinase, also called PIK3CA, is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. PIK3CA gene is mapped to 3q26.32. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate phosphatidylinositols (PtdIns), PtdIns4P and PtdIns(4,5)P₂. Recent evidence has shown that the PIK3CA gene is mutated in a range of human cancers. It has been found to be oncogenic and has been implicated in cervical cancers. PIK3CA mutations in breast cancer may be a predictive marker to guide the selection of patients who would benefit from mTOR inhibitor therapy. In addition to that, the presence of PIK3CA mutation may predict response to aspirin therapy for colorectal cancer, indicating power and promise of "Molecular Pathological Epidemiology (MPE)" approach as well as a complex interaction within the tumor microenvironment in this phenomenon.

Reference

Anti-PI3 Kinase p110 Alpha/PIK3CA Antibody被引用在2文献中。

Selected Validation Data




Figure 1. Western blot analysis of PIK3CA using anti-PIK3CA antibody (PB0351). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: SW620 Whole Cell Lysate, Lane 2: PC12 Whole Cell Lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PIK3CA antigen affinity purified polyclonal antibody (Catalog # PB0351) at 0.5 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PIK3CA at approximately 110KD. The expected band size for PIK3CA is at 110KD.