

Basic Information

Product Name	Anti-Prolactin/PRL Antibody
Gene Name	PRL
Source	Rabbit
Isotype	IgG
Species Reactivity	human, rat
Tested Application	WB, ELISA
Contents	500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol.
Immunogen	E.coli-derived human Prolactin recombinant protein (Position: L29-227C). Human Prolactin shares 60% and 63% amino acid (aa) sequence identity with mouse and rat Prolactin, respectively.
concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	26KD
Dilution Ratios	Western blot(WB):1:500-2000 ELISA: 1:100-1000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

Prolactin (PRL), also known as luteotropic hormone or luteotropin, is a protein that in humans is best known for its role in enabling female mammals to produce milk. This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants that encode the same protein.

Reference

Anti-Prolactin/PRL Antibody被引用在2文献中。

Selected Validation Data



Figure 1. Western blot analysis of Prolactin using anti-Prolactin antibody (PB0364). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: Rat Brain Tissue Lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Prolactin antigen affinity purified polyclonal antibody (Catalog # PB0364) at 0.5 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Prolactin at approximately 26KD. The expected band size for Prolactin is at 26KD.