

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Product Name Anti-TSG101 Antibody Gene Name TSG101 Source Rabbit Isotype IgG Species Reactivity human, rat Tested Application WB Contents 500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol. Immunogen A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD Dilution Ratios Western blot(WB):1:500-2000	Basic Information	
Source Rabbit Isotype IgG Species Reactivity human, rat Tested Application WB Contents 500 ug/ml antibody with PBS + 0.02% NaN3 + 1 mg BSA and 50% glycerol. Immunogen A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD	Product Name	Anti-TSG101 Antibody
Isotype IgG	Gene Name	TSG101
Species Reactivity human, rat Tested Application WB Contents 500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol. Immunogen A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD	Source	Rabbit
Tested Application Contents 500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol. A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD	Isotype	IgG
Contents 500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol. Immunogen A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD	Species Reactivity	human, rat
Immunogen A synthetic peptide corresponding to a sequence at the C-terminus of human TSG101 (361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. 0bserved MW 44KD	Tested Application	WB
(361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and sequences. concentration 500 ug/ml Purification Immunogen affinity purified. Observed MW 44KD	Contents	500 ug/ml antibody with PBS $_{\odot}$ 0.02% NaN3 , 1 mg BSA and 50% glycerol.
Purification Immunogen affinity purified. Observed MW 44KD	Immunogen	(361-390aa KHVRLLSRKQFQLRALMQKARKTAGLSDLY), identical to the related mouse and rat
Observed MW 44KD	concentration	500 ug/ml
	Purification	Immunogen affinity purified.
Dilution Ratios Western blot(WB):1:500-2000	Observed MW	44KD
	Dilution Ratios	Western blot(WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

TSG101, known as Tumor susceptibility gene 101, is mapped to 11p15. The protein encoded by this gene belongs to a group of apparently inactive homologs of ubiquitin-conjugating enzymes. The gene product contains a coiled-coil domain that interacts with stathmin, a cytosolic phosphoprotein implicated in tumorigenesis. And the protein may play a role in cell growth and differentiation and act as a negative growth regulator. In vitro steady-state expression of this tumor susceptibility gene appears to be important for maintenance of genomic stability and cell cycle regulation. Mutations and alternative splicing in this gene occur in high frequency in breast cancer and suggest that defects occur during breast cancer tumorigenesis and/or progression.

Reference

Anti-TSG101 Antibody被引用在1文献中。

BOSTER BIOLOGICAL TECHNOLOGY

Special NO.1, International Enterprise Center, 2nd Guanshan Road, Wuhan, China

Web: www.boster.com.cn Phone: +86 027-67845390 Fax: +86 027-67845390 Email: boster@boster.com.cn

Selected Validation Data




Figure 1. Western blot analysis of Anti-TSG101 antibody (PB0550). The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: Rat cardiac muscle tissue lysates,Lane 2: Rat brain tissue lysates,Lane 3: HELA whole cell lysates,Lane 4: SMMC whole cell lysates,Use rabbit Anti-TSG101 1:1000, probed with a goat Anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002). A specific band was detected for TSG101 at approximately 44KD. The expected band size for TSG101 is at 44KD.