

Basic Information

Product Name	Anti-HO-1/HMOX1 Antibody	
Gene Name	HMOX1	
Source	Rabbit	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol.	
Immunogen	E.coli-derived human HMOX1 recombinant protein (Position: M1-M288). Human HMOX1 shares 82% and 80% amino acid (aa) sequences identity with mouse and rat HMOX1, respectively.	
concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	33KD	
Dilution Ratios	Western blot(WB): 1:500-2000 Immunohistochemistry in paraffin section (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.

Background Information

HMOX1 (heme oxygenase (decycling) 1), also known as HO-1, is a human gene that encodes for the enzyme heme oxygenase 1. It is an essential enzyme in heme catabolism, it cleaves heme to form biliverdin. HMOX1 belongs to the heme oxygenase family. The HMOX1 gene is located on the long (q) arm of chromosome 22 at position 12.3, from base pair 34,101,636 to base pair 34,114,748. HMOX1, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is subsequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitter. HMOX1 activity is induced by its substrate heme and by various nonheme substances.

Reference

Anti-HO-1/HMOX1 Antibody被引用在4文献中。

Selected Validation Data




Figure 1. Western blot analysis of anti-HMOX1 antibody (PB9212).

The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: A549 Whole Cell Lysate,

Lane 2: HELA Whole Cell Lysate.

Use rabbit anti-HMOX1 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002). A specific band was detected for HMOX1 at approximately 33KD. The expected band size for HMOX1 is at 33KD.




Figure 2. IHC analysis of HMOX1 antibody (PB9212). was detected in paraffin-embedded section of human tonsil tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.