

Basic Information

Product Name	Anti-Survivin/BIRC5 Antibody	
Gene Name	BIRC5	
Source	Rabbit	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB,IHC,ELISA	
Contents	500 ug/ml antibody with PBS , 0.02% NaN3 , 1 mg BSA and 50% glycerol.	
Immunogen	E. coli-derived human Survivin recombinant protein(Position: M1-D142).	
concentration	500 ug/ml	
Observed MW	16-18KD	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry in paraffin section (IHC): 1:50-400 ELISA 1:10-1000 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied. 6 months 2 to 8°C after reconstitution. Avoid repeated freezing and thawing.


Background Information

Survivin, also called baculoviral inhibitor of apoptosis repeat-containing 5 or BIRC5, is a protein that, in humans, is encoded by the BIRC5 gene. Survivin is a member of the inhibitor of apoptosis(IAP) family. The survivin gene contains 4 exons. The survivin gene is mapped to chromosome 17q25 by pulsed field gel electrophoresis and single- and 2-color FISH. The survivin protein functions to inhibit caspase activation, thereby leading to negative regulation of apoptosis or programmed cell death. The survivin protein is expressed highly in most human tumours and fetal tissue, but is completely absent in terminally differentiated cells.

Reference

Anti-Survivin/BIRC5 Antibody被引用在2文献中。

Selected Validation Data



Anti-human Survivin antibody, RP1026, Western blotting Lane 1:
Recombinant Human Survivin Protein 10ng Lane 2: Recombinant
Human Survivin Protein 5ng Lane 3: Recombinant Human Survivin
Protein 2